



*In Partnership with* **CalViva**  
HEALTH

# Coding for Vascular Conditions

# Agenda

Vascular Diseases

The Impact on Risk Adjustment Payment Model

ICD-10-CM Guidance

Coding Trends

The Role of the Provider

Resources

---

# Vascular Diseases

---



# Vascular Disease



Affects circulatory system which ranges from diseases of arteries, veins and lymph vessels to blood disorders that affect circulation.



Smoking and Diabetes are common risk factors.



Restricts the amount of blood that circulates throughout the vessels of the extremities. With this, your skin and tissues in the extremities receive less oxygen and vital nutrients to stay healthy and heal.



Over time, the main risks are stroke, heart attack, pulmonary embolism, and death.



In Partnership with  CalViva Health

# Vascular Disease

Circulatory System  
Arterial and Venous



Photo credit: [www.cdc.gov/TSp/substances/ToxOrganListing.aspx?toxid=1](http://www.cdc.gov/TSp/substances/ToxOrganListing.aspx?toxid=1)

## Restricted blood flow can cause:

- Inflammation – narrowing of the vessels from swelling
- Claudication – weakness, leg pain, heaviness and cramping of the muscles. Happens when active and stops (or diminishes) when at rest
- Arteriosclerosis/Atherosclerosis – fatty deposits blocking blood vessels
- Thrombus – a clot/blockage of the blood vessel
- Ulcers – sores that heal slowly or won't heal

## Prevention:

- Don't smoke
- Control blood sugar
- Exercise regularly
- Maintain healthy weight
- Manage blood pressure

# Peripheral Vascular Disease I73.9

- Includes:
  - Intermittent claudication
  - Peripheral angiopathy NOS
  - Spasm of artery
- Contains an Excludes1 Note
  - Atherosclerosis of the extremities (I70.2-I70.7-)

## ICD-10-CM Tabular List

### I73.9 Peripheral vascular disease, unspecified

Intermittent claudication  
Peripheral angiopathy NOS  
Spasm of artery

**EXCLUDES1** atherosclerosis of the extremities (I70.2-I70.7-)

# Atherosclerosis I70.xxx






Photo credit:  
<https://www.nhlbi.nih.gov/health/atherosclerosis>

# Venous Ulcers

- Venous ulcers are a common side effect of PVD.
- Do not use 'wound' if it is an ulcer
  - Wound is an injury that happens traumatically
  - Ulcer is a breakdown of the skin that fails to heal
- Documentation is key to ulcer coding:
  - Type
  - Site
  - Laterality
  - Severity
  - Underlying condition





# Arteriosclerosis of Arteries of Extremities with Complication and Ulceration

## Documentation Tips

Documentation for arteriosclerosis of lower extremities should include:

- ✓ **System:** Arterial or venous
- ✓ **Vessel type:** Native or graft (autologous, non-autologous biologic, non-autologous nonbiologic)
- ✓ **Anatomic location:** Include site, laterality
- ✓ **Complicating factors:** Intermittent claudication, rest pain, ulceration, and gangrene
  - ✓ Use additional documentation to report **the non-pressure ulcer (L97.-)**, if applicable, by documenting :
    - Site
    - Laterality
    - Depth
- ✓ “Healing” ulcers are considered as active
- ✓ “Healed” ulcers are considered resolved

## Common Tests / Treatment


### Tests

- ✓ Pressure measurements: Ankle-brachial index
- ✓ Exercise testing
- ✓ Toe systolic pressure index
- ✓ Segmental pressures and pulse volume recordings
- ✓ Flow velocity determination
- ✓ Ultrasonic duplex scanning

### Treatment

- ✓ Exercise therapy (intermittent claudication)
- ✓ Smoking cessation
- ✓ Drug therapy (vasodilators, rheologic agents, anhydrobiotic therapy)
- ✓ Interventional procedures
- ✓ Wound care

# Skin Ulcers – Specificity and Combination Codes



**Atherosclerosis of lower extremities**  
 Ex. I70.244 Atherosclerosis of native arteries of left leg **with** ulcer of left heel and midfoot (Hierarchical Condition Category (HCC) 161 & HCC 106) (HCC 263 AND HCC 383) L97.429 Non-pressure chronic ulcer of left heel of unspecified severity.

**Diabetes**  
 Ex. E11.621 Type 2 diabetes mellitus **with** foot ulcer (HCC1 8 & HCC 161) (37 AND 383)

**Chronic venous hypertension**  
 Ex. I87.332 Chronic venous hypertension **with** ulcer and inflammation of left lower extremity (HCC 107) ( HCC 383)

**Post-thrombotic syndrome**  
 Ex. I87.032 Post-thrombotic syndrome **with** ulcer and inflammation of the left lower extremity (HCC 107) (HCC 383)

**Varicose ulcer**  
 Ex. I83.024 Varicose veins of left lower extremity **with** ulcer of heel and midfoot (HCC 107) ( HCC 383)

**Any associated gangrene**  
 Ex. I70.262 Atherosclerosis of native arteries of extremities **with** gangrene (and ulcer), left leg (HCC 106) (HCC 263)



# Aneurysms I71.xxx and I72.X




Photo credit: [www.nhlbi.nih.gov/health/aortic-aneurysm](http://www.nhlbi.nih.gov/health/aortic-aneurysm)

Aneurysms are an abnormal bulge in the wall of a blood vessel.

They are caused by atherosclerosis, high blood pressure, injury to your aorta, and family history of aneurysms.

Depending on the type, size and shape treatment can include medications or surgery.

Unruptured aneurysms are monitored closely with the goal of treatment to prevent the aneurysm from bursting.

# Aneurysms

## Documentation / coding tips

### Coding tips:

- Use the current year of ICD-10-CM book (updates every October 1) for accurate coding assignment.
- Carefully review all codes first, excludes 1 and 2 notes in the tabular list before selecting a code.

### Documentation tips:

- Must specify the type, location, size, whether rupture has occurred, and if it has been repaired.
  - History of Surgical Repair – Z86.79
  - Aortic (Aorta) Aneurysm – I71.xx
  - Other Aneurysm – I72.xx (include laterality as applicable)
- Detail is needed. Coders can not make assumptions. Documentation must be clear, concise, and complete.

# Embolism and Thrombosis





Photo credit: [www.cdc.gov/blood-clots/index.html](http://www.cdc.gov/blood-clots/index.html)

# Embolism and Thrombosis

## Documentation / coding tips

### Coding tips:

- Thrombosis happens when a blood clot grows in blood vessels that reduce blood flow.
- Embolus is any foreign material that travels and becomes stuck which blocks the flow of blood.
- Vascular occlusion is any term of blockage of blood vessel.

### Documentation tips:

- Must specify the type, location, and any complications, and whether it caused any complications such as stroke.
  - Aorta abdominal thrombosis I74.09
  - Carotid (artery) thrombosis see occlusion, artery, carotid I65.2X
  - Upper extremity embolism – I74.2
  - Series I63.XXX cerebral infarction due to thrombosis, embolism, and occlusion
  - Past history of venous embolism or thrombosis Z86.718

# Deep Vein Thrombosis

## Documentation Tips

- ✓ Distinguish between **DVT** and **venous thrombosis with phlebitis**
- ✓ Clearly document whether DVT is **acute, chronic or historical**
- ✓ The **treatment and its goals** should be linked to the diagnosis
- ✓ The documentation should indicate:
  - ✓ **Laterality** – side of the body, i.e., left or right
  - ✓ The **specific location** – upper or lower extremity
  - ✓ The **specific vein** such as femoral, iliac or tibial

**Note:** A coder cannot assume a DVT is acute, chronic or historical based on suggestive documentation or on a timeline of events. The condition can only be coded based on documentation.

## Common Tests and Treatment


### Tests

- ✓ Venous ultrasound
- ✓ Venography
- ✓ D-Dimer blood test
- ✓ Impedance plethysmography
- ✓ Magnetic resonance imaging (MRI)

### Treatment

- ✓ Anticoagulant medications
- ✓ Inferior vena cava (IVC) filter
- ✓ Compression stockings

# Deep Vein Thrombosis (DVT) – Examples:






# Cerebrovascular Accident (CVA)

## Coding for CVA status post discharge

### Inpatient setting



**I63.xxx – Cerebral infarction**  
*Acute CVA should only be documented during the initial episode of care.*

### Outpatient setting

Residual deficit(s)?

Yes


#### **I69.3xx – Sequelae of cerebral infarction**

- Report sequelae for as long as they are present.
  - Cognitive deficit
  - Speech/language deficit
  - Monoplegia/Hemiplegia/Hemiparesis
- Use as many codes from category I69.3 as needed to identify all late effects.
- Documentation should clearly tie deficit to CVA:
  - “due to CVA”*
  - “late effect of CVA”*

No

**Z86.73 - Personal history of transient ischemic attack (TIA), and CVA without residual deficits**

# Sequelae of CVA (Stroke) – Category I69



## I69 - Sequelae of cerebrovascular disease

- Conditions that are specified as “*sequelae*” or as “*residuals*” of cerebrovascular disease should be reported using a code from category I69
- Sequelae may arise at any time after the onset of the cerebrovascular condition
- Documentation should clearly link sequela(e) to cerebrovascular condition using language such as:
  - “*sequela of CVA*”
  - “*residual of CVA*”
  - “*late effect of CVA*”
  - “*associated with CVA*”
  - “*due to CVA*”

### Unilateral weakness caused by CVA

When unilateral weakness is clearly documented as being associated with a stroke, it is considered synonymous with **hemiparesis/hemiplegia**. Unilateral weakness outside of this clear association cannot be assumed as hemiparesis. (*Coding Clinic, 2015, 1Q, 25*)

### Upper or lower extremity weakness following CVA

Weakness of one extremity (upper or lower) that is clearly associated with CVA should be reported as **monoplegia**. (*Coding Clinic, 2017, 1Q, 47*)

### Dominant vs. nondominant side

Codes from category I69 specify whether the dominant or non-dominant side is affected. Should the affected side (i.e., right, left) be documented but not identified as dominant or non-dominant, code selection is as follows:

- If the right side is affected, the default is dominant.
- If the left side is affected, the default is non-dominant.
- For ambidextrous patients, the default should be dominant.

---

# Vascular Conditions and the Impact on Risk Adjustment Payment Model

---


---

# Risk Adjustment Overview

---

# What is Risk Adjustment?

The Centers for Medicare & Medicaid Services (CMS) Risk Adjustment Program uses a predictive algorithm that incorporates information on individuals' demographics and health conditions to predict variation in future medical expenditures. The model is used to forecast trends and future needs of patients.



Accurate risk adjustment relies on complete documentation and coding to the highest level of specificity known.

# Why is Risk Adjustment Important?



Accuracy in member health status profile



Improved quality of care through disease management programs



Appropriate risk premium

---

# Risk Adjustment HCC CMS Mapping Changes (V28)

---

# High Level Overview of Medicare Advantage (MA) Model Changes

THREE YEAR PHASE IN		
Payment Year	RA Model (v24)	RA Model (v28)
PY2024 (DOS 2023)	67%	33%
PY2025 (DOS 2024)	33%	67%
PY2026 (DOS 2025)	0%	100%

✓ Familiar diagnosis at low specificity deleted across disease categories

Example: Angina pectoris

✓ New codes added to disease categories

Example: J45.51 – Severe persistent asthma

Example: F50.00 – Anorexia nervosa

✓ HCCs added

Example: HCC 279 – Severe persistent asthma


Example: HCC 153 – Anorexia/bulimia nervosa

✓ HCC numbers and coefficient update

Example: Diabetes w/comp

HCC18 to HCC37

0.318 to 0.166





# V24 vs V28 HCC Categories

Description	ICD-10	V24	V28
Acute embolism and thrombosis of unspecified deep veins of right lower extremity	I82.401	108	267
Acute embolism and thrombosis of unspecified deep veins of left lower extremity	I82.402	108	267
Chronic embolism and thrombosis of unspecified deep veins of right lower extremity	I82.501	108	267
Chronic embolism and thrombosis of unspecified deep veins of left lower extremity	I82.502	108	267
Atherosclerosis of extremities (native arteries) with rest pain, left leg	I70.222	108	264
Atherosclerosis of extremities (native arteries) with rest pain, right leg	I70.221	108	264
Varicose veins of right lower extremity ulcer of ankle	I83.013	107	383
Varicose veins of right lower extremity ulcer of heel and midfoot	I83.014	107	383

# V24 vs. V28 Vascular Disease

V24	V28
<ul style="list-style-type: none"><li>• Vascular disease group:<ul style="list-style-type: none"><li>• <b>HCC 106</b> Atherosclerosis of the extremities with ulceration or gangrene</li><li>• <b>HCC 107</b> Vascular disease with complications</li><li>• <b>HCC 108</b> Vascular disease</li></ul></li></ul>	<ul style="list-style-type: none"><li>• Vascular disease group:<ul style="list-style-type: none"><li>• <b>HCC 263</b> Atherosclerosis of arteries of the extremities with ulceration or gangrene</li><li>• <b>HCC 264</b> Vascular disease with complications</li><li>• <b>HCC 267</b> Deep vein thrombosis and pulmonary embolism</li></ul></li></ul>

## Dx removed from v28 Model:

- Atherosclerosis of aorta
- Atherosclerosis of renal artery
- Atherosclerosis of native arteries of extremities with intermittent claudication
- Aortic aneurysm, without rupture
- Aortic ectasia
- Peripheral vascular disease

- ❖ Only atherosclerosis with complications remain on the model.

## Documentation best practice

- ❖ Complications of atherosclerosis of arteries should be documented and supported in the medical record when they are present

---

# ICD-10 Coding Guidelines

---

# Excludes Notes – Excludes1

## Excludes1



- Means “**NOT CODED HERE!**”
- Indicates two conditions cannot occur together.

*Example*

✓4<sup>th</sup>

**I82.7 Chronic embolism and thrombosis of veins of upper extremity**

**Use additional code, if applicable , for associated long- term (current) use of anticoagulants (Z79.01)**

**EXCLUDES 1**

*personal history of venous embolism and thrombosis (Z86.718)*

*Cannot report Chronic embolism and thrombosis of veins of upper extremity with personal history of venous embolism and thrombosis*

# Excludes Notes – Excludes2

**Excludes2**

- Means “**Not included here**”
- Excluded condition is not represented by the code above
- Patient may have both conditions

**✓4<sup>th</sup> I83 Varicose veins of lower extremities**

**EXCLUDES 2** *Varicose veins complicating pregnancy (O22.0-)*  
*Varicose veins complicating puerperium (O87.4)*

Category I83 does not include Varicose veins in pregnancy or puerperium.

ICD-10 Code

Included Condition(s)

Excluded Condition

# Section IV. Diagnostic Coding and Reporting Guidelines for Outpatient Services

## D. Codes that describe symptoms and signs

Codes that describe symptoms and signs, as opposed to diagnoses, are acceptable for reporting purposes when a diagnosis has not been established (confirmed) by the physician or other provider. Chapter 18 of ICD-10-CM, Symptoms, Signs, and Abnormal Clinical and Laboratory Findings Not Elsewhere Classified (codes R00-R99) contain many, but not all codes for symptoms.

## G. ICD-10-CM code for the diagnosis, condition, problem, or other reason for encounter/visit

List first the ICD-10-CM code for the diagnosis, condition, problem, or other reason for encounter/visit shown in the medical record to be chiefly responsible for the services provided. List additional codes that describe any coexisting conditions. In some cases, the first-listed diagnosis may be a symptom when a diagnosis has not been established (confirmed) by the physician or other provider.

## H. Uncertain diagnosis

Do not code diagnoses documented as “probable,” “suspected,” “questionable,” “rule out,” “compatible with,” “consistent with,” or “working diagnosis,” or other similar terms indicating uncertainty. Rather, code the condition(s) to the highest degree of certainty for that encounter/visit, such as symptoms, signs, abnormal test results, or other reason for the visit.

**Please note:** This differs from the coding practices used by short-term, acute care, long-term care and psychiatric hospitals.

# Section I. Conventions, general coding guidelines and chapter specific guidelines

## 10. Includes Notes

This note appears immediately under a three-character code title to further define, or give examples of, the content of the category.

## Example

### I74 Arterial embolism and thrombosis

#### INCLUDES

embolic infarction  
embolic occlusion  
thrombotic infarction  
thrombotic occlusion

## 11. Inclusion terms

List of terms is included under some codes. These terms are the conditions for which that code is to be used. The terms may be synonyms of the code title, or, in the case of “other specified” codes, the terms are a list of the various conditions assigned to that code. The inclusion terms are not necessarily exhaustive. Additional terms found only in the Alphabetic Index may also be assigned to a code.

## Example

### I73.9 Peripheral vascular disease, unspecified

Intermittent claudication  
Peripheral angiopathy NOS  
Spasm of artery

# ICD 10-CM Convention Guideline

Per the ICD-10 Official Guidelines, the word “with” or “in” should be interpreted to mean “associated with” or “due to” when it appears in a code title, the Alphabetic Index, or an instructional note in the Tabular List. The classification presumes a causal relationship between the two conditions linked by these terms in the Alphabetic Index or Tabular List. These conditions should be coded as related even in the absence of physician or other provider documentation explicitly linking them, unless the documentation clearly states the conditions are unrelated or when another guideline exists that specifically requires a documented linkage between two conditions.

Diabetes, diabetic (mellitus) (sugar) E11.9

- with

- chronic kidney disease E11.22
- circulatory complication NEC E11.59
- coma due to
- hyperosmolarity E11.01
- hypoglycemia E11.641
- ketoacidosis E11.11
- complication E11.8
- specified NEC E11.69
- dermatitis E11.620
- foot ulcer E11.621
- gangrene E11.52
- gastroparesis E11.43
- gastroparesis E11.43
- glomerulonephrosis, intracapillary E11.21
- glomerulosclerosis, intracapillary E11.21
- hyperglycemia E11.65
- hyperosmolarity E11.00
- with coma E11.01
- hypoglycemia E11.649
- with coma E11.641
- ketoacidosis E11.10
- with coma E11.11
- kidney complications NEC E11.29
- ~~Wilson disease E11.21~~
- loss of protective sensation (LOPS) -see Diabetes, by type, with neuropathy
- mononeuropathy E11.41
- myasthenia E11.44
- necrobiosis lipoidica E11.620
- nephropathy E11.21
- neuralgia E11.42
- neurologic complication NEC E11.49
- neuropathic arthropathy E11.610
- neuropathy E11.40
- ophthalmic complication NEC E11.39
- oral complication NEC E11.638
- osteomyelitis E11.69
- periodontal disease E11.630
- peripheral angiopathy E11.51
- with gangrene E11.52

*These conditions should be coded as related to diabetes, even in the absence of physician or other provider documentation explicitly linking them.*

**Coding tip** - A causal relationship between diabetes and above conditions is reported UNLESS:

- Documentation identifies another cause for the condition(s)
- Documentation clearly states the condition(s) are not caused by diabetes
- Documentation explicitly states underlying cause of condition(s) is unknown, under workup, etc.
- Any condition with NEC (not elsewhere classifiable) following it requires the physician or other provider to document the relationship



In Partnership with CalViva Health



---

# Coding Trends

---

# Example – Specificity

## Past Medical History

Reviewed history from 01/13/2023 and no changes required:

Diabetes, Type 2

Hypertension

Legally blind

PVD post amputation

Clearance for cataract surgery

## Impression & Recommendations:

**Problem #1:** Diabetes, Type 2 (ICD9 - 250.00) (ICD10 - E11.9)

**Assessment:** Unchanged.

His updated medication list for this problem includes:

Tresiba FlexTouch U-100 100 Unit/ml (3M) Insulin Pen (Insulin degludec)

...Inject 50 units subcutaneously once a day.

Irbesartan-hydrochlorothiazide 300-12.5 Mg Tablet (Irbesartan-hydrochlorothiazide) ...1 tablet by mouth once a day

Glyburide-metformin 5-500 Mg tablet (Glyburide-metformin) ... Take 1 tablet by mouth twice a day.

## ICD-10-CM Guidelines

### 15. “With”

The word “with” or “in” should be interpreted to mean “associated with” or “due to” when it appears in a code title, the Alphabetic Index (either under a main term or subterm), or an instructional note in the Tabular List. The classification presumes a causal relationship between the two conditions linked by these terms in the Alphabetic Index or Tabular List. These conditions should be coded as related even in the absence of physician or other provider documentation explicitly linking them, unless the documentation clearly states the conditions are unrelated or when another guideline exists that specifically requires a documented linkage between two conditions.

## ICD-10-CM Alphabetic Index

type 2 E11.9

With

osteomyelitis E11.69

periodontal disease E11.630

peripheral angiopathy E11.51

polyneuropathy E11.42

In this example, Type 2 diabetes E11.9 is not coded to the highest level of specificity. This is due to a presumed causal relationship between diabetes and PVD.

**Correct code:**

E11.51 Type 2 diabetes mellitus with peripheral vascular disease

# Example - Missed Opportunities

## Physical examination:

**Skin:** Inspection and palpation: no rash, ulcer, induration, nodules, jaundice, or abnormal nevi and good turgor and lesion (dressing in place of left heel – stasis ulcer picture from daughter).

## 2. Venous stasis ulcer of leg – Left

I83.009: *Varicose veins of unspecified lower extremity with ulcer of unspecified site*

I83.023: *Varicose veins of left extremity with ulcer of ankle*

WOUND CARE REFERRAL – Schedule within provider's discretion.

Note to provider: patient has venous stasis ulcer to left heel/ankle; would like home health once set up with you

In this example, the varicose veins ulcer of the leg was in assessment with plan treatment documented. Under I83.0 series in the tabular list, use additional code to identify severity of the ulcer L97.-

### Correct code:

Heel was in exam, and ankle and heel was in assessment  
I83.023 (ankle) I83.024 (heel)  
L97.329 non-pressure ulcer left ankle  
unspecified severity  
L97.429 non- pressure ulcer left heel  
unspecified severity

# Examples - Diagnosis Not Supported

## Assessments:

1. Well adult exam – Z00.00 (Primary)
2. Constipated – K59.00
3. Acute deep vein thrombosis (DVT) of distal vein of left lower extremity – I82.4Z2

## Plan:

Acute deep vein thrombosis (DVT) of distal vein of left lower extremity.

Refill Xarelto tablet, 20mg, 1 tab(s), orally, once a day (in the evening), 30 day(s), 30, Refills 3

Notes: Med adherence stressed.

## Past Medical History

HTN.

History of L extremity DVT.

History of bilateral popliteal DVT.

Severe lumbar stenosis.

Internal hemorrhoids.

In this example, there is not enough support to show this acute condition is active. Proof of the clot and/or date condition was first diagnosed and treatment started could support this. History of acute DVT is also documented in past medical history could be conflicting documentation.

# Examples - Diagnosis Not Supported

## Chief Complaint

- Comprehensive health assessment
- Follow-up on chronic conditions
- Provided printed screening plan to patient.
- Follow up on second degree AV block.

68 y/o male presents for WRV.

### #Nonsustained supraventricular tachycardia / second degree AV block:

-Pt. denies cp today. Occasionally he has palpitations. He has not had any recent follow ups with cardiology and has not been taking any of his medications.

### #Hypertensive heart disease:

-Pt. reports non-compliance with all his medications.

-Pt. reports feeling dizzy and lightheaded when he goes from sitting to standing. Pt. denies exhaustion.

### #T2DM:

-Pt. has Lantus at home but hasn't been taking it.

-He reports his pharmacy didn't have any pen needles when picked up his medications.

### Assessment:

- R/O PAD (peripheral vascular disease) 443.9/I73.9 5/24/2023.
- Per QF 5/10/23: left foot = 0.98 (Mild)
- Checking lipids with labs today
- T/c starting statin

## ICD-10-CM Guidelines

### Section IV. Diagnostic Coding and Reporting Guidelines for Outpatient Services

#### Uncertain diagnosis

Do not code diagnoses documented as “probable”, “suspected,” “questionable,” “rule out,” “compatible with,” “consistent with,” or “working diagnosis” or other similar terms indicating uncertainty. Rather, code the condition(s) to the highest degree of certainty for that encounter/visit, such as symptoms, signs, abnormal test results, or other reason for the visit.

**Please note:** This differs from the coding practices used by short-term, acute care, long-term care and psychiatric hospitals.

In this example, Rule out (R/O) PAD was documented in the assessment and captured as I73.9 Peripheral artery disease (PAD).

Per coding guidelines, do not code diagnoses documented as “rule out” or other similar terms indicating uncertainty.



---


# The Role of the Provider

---

# Best Practices – Documentation Strategies

Clear, Concise, Complete

***Documentation should paint a picture with words of the patient's condition and what occurred during each visit.***



# What Can You Do?

The two most important things that physicians and other providers can do regarding risk adjustment are:

See the patient at least once a year to determine health status.

- Evaluate and document ALL active conditions.
- Simply listing every diagnosis in the medical record is not acceptable and does not support reporting an HCC.

Be as specific as possible in the documentation.

- This will allow for the most accurate ICD-10-CM code to be reported.
- Documentation should include any manifestations or complications related to a chronic disease.



# “History of” vs. Active Condition

## HISTORY OF

- Medical conditions that no longer exist or have resolved should not be reported as active.
- History codes are used to explain a patient’s past medical condition for which they are no longer receiving active treatment.

## ACTIVE

- Medical conditions can occur suddenly and last a short period of time, such as a few days or weeks.
- An acute condition should be coded when present and actively being treated.
- Medical record documentation needs to support the active/acute condition.

# Key Takeaways

Reviewing coding guidelines annually - chart reviews of high-risk diagnoses to assure compliance with guidelines.

Use both alphabetic index and tabular list when selecting codes will ensure all excludes 1 & 2, code first, and use additional codes notes are reviewed.

Code to the highest degree of specificity and ensure the diagnoses are properly sequenced on the claim.

Conditions that were previously treated and no longer exist should be coded using history of codes.

---

# Resources

---

# What can the CDI team do for you

## Our Mission

- ❑ Help physicians and other providers understand and apply risk adjustment concepts. We can also help in the application of documentation and coding best practices to workflows.
- ❑ Protect the integrity and accuracy of risk adjusted diagnosis and improve outcomes.



## Our Services

- ❑ Live Webinars
  - Risk Adjustment Concepts 101
  - RA Documentation & Coding Best Practices
  - Avoiding Common Documentation Errors
  - Yearly ICD-10-CM Updates
  - Disease Specific
- ❑ Train the Trainer
  - 1:1 Coder training
  - RA Documentation Requirements
- ❑ Chart reviews
  - Baseline reviews
    - Onboarding for new physicians and other providers
    - General assessment for practice/provider
  - Targeted review
    - Disease/HCC/ICD-10 specific
  - Tailored education based on chart review findings
    - Share identified opportunities to provide greater specificity to coding and documentation
    - Address gaps in documentation
- ❑ Disease-Specific Coding Reference Material
  - human immunodeficiency virus, diabetes mellitus, congestive heart failure and many more

## The Goal

- ❑ Engage staff and entire team in learning
- ❑ Increase HCC proficiency
- ❑ Enhanced communication support between coding staff, administrative staff, physicians, and other providers
- ❑ Increase awareness of implications related to inaccurate coding

Services are complimentary.  
For more information,  
please email  
[askcdi@centene.com](mailto:askcdi@centene.com)

# For Requests and Questions



[AskCDI@centene.com](mailto:AskCDI@centene.com)

Education requests, coding and documentation questions, etc.



[CDIwebinars@centene.com](mailto:CDIwebinars@centene.com)

CDI webinar requests for providers and health plans

- Risk Adjustment 101 – All LOBs
- Diagnosis/HCC specific training
- Claims coding/billing training – specific to claims denials/rejects regarding HCC related diagnosis codes



[RAPS@centene.com](mailto:RAPS@centene.com)

Provider can email records to CDI educator or fax records to our secure fax: 813-464-8900

---

# Thank you for your attendance!

---